

Table 11.1. Mean catch-per-unit-effort and (standard error) for fishes collected by large hoop netting in Pool 4 of the Mississippi River using stratified random sampling in 2007. The statistics under ALL pertain to unbiased means over ALL strata sampled by this gear (as indicated by nonmissing entries below and by Table 2.1). See methods for definitions of catch-per-unit-effort and standard error.

Common name	ALL	Main channel border, unstructured	Side channel border
Bowfin	0.02 (0.02)		0.04 (0.04)
Gizzard shad	0.03 (0.03)	0.06 (0.06)	
Common carp	0.87 (0.53)	0.19 (0.13)	1.39 (0.93)
River carpsucker	0.09 (0.09)	0.21 (0.21)	
Smallmouth buffalo	1.39 (0.79)	2.28 (1.78)	0.70 (0.23)
Black buffalo	0.02 (0.02)		0.04 (0.04)
Spotted sucker	0.02 (0.02)		0.04 (0.04)
Silver redhorse	0.18 (0.09)	0.20 (0.14)	0.17 (0.11)
River redhorse	0.02 (0.02)		0.04 (0.04)
Golden redhorse	0.11 (0.11)	0.25 (0.25)	
Shorthead redhorse	0.30 (0.16)	0.48 (0.34)	0.16 (0.09)
Channel catfish	1.82 (0.69)	2.91 (1.40)	0.97 (0.58)
Flathead catfish	0.18 (0.11)	0.19 (0.13)	0.17 (0.17)
Northern pike	0.21 (0.08)	0.13 (0.08)	0.28 (0.14)
White bass	0.42 (0.17)	0.25 (0.13)	0.56 (0.28)

Table 11.1 continued.

Common name	ALL	Main channel border, unstructured	Side channel border
Rock bass	0.03 (0.03)	0.06 (0.06)	
Bluegill	1.01 (0.40)	0.70 (0.49)	1.25 (0.60)
Largemouth bass	0.02 (0.02)		0.04 (0.04)
White crappie	0.07 (0.03)		0.12 (0.06)
Black crappie	1.44 (0.60)	2.09 (1.21)	0.94 (0.50)
Walleye	0.08 (0.04)	0.13 (0.09)	0.04 (0.04)
Freshwater drum	1.79 (0.57)	2.82 (1.20)	0.98 (0.39)