

Table 6.3. Mean catch-per-unit-effort and (standard error) for fishes collected by fyke netting in Pool 13 of the Mississippi River using stratified random sampling in 2007. The statistics under ALL pertain to unbiased means over ALL strata sampled by this gear (as indicated by nonmissing entries below and by Table 2.3). See methods for definitions of catch-per-unit-effort and standard error.

Common name	ALL	Backwater, contiguous, shoreline	Impounded, shoreline
Longnose gar	0.06 (0.05)	0.05 (0.05)	0.13 (0.13)
Shortnose gar	0.62 (0.40)	0.64 (0.44)	0.49 (0.32)
Bowfin	0.88 (0.58)	0.98 (0.65)	
Gizzard shad	8.50 (3.00)	9.26 (3.32)	1.27 (0.79)
Common carp	2.80 (0.83)	2.82 (0.90)	2.62 (1.49)
Golden shiner	2.24 (1.42)	2.42 (1.57)	0.52 (0.52)
River carpsucker	0.20 (0.11)	0.20 (0.12)	0.13 (0.13)
Highfin carpsucker	0.07 (0.06)		0.75 (0.61)
Spotted sucker	0.14 (0.08)	0.16 (0.09)	
Shorthead redhorse	0.97 (0.72)	1.06 (0.80)	0.12 (0.12)
Yellow bullhead	0.44 (0.34)	0.48 (0.38)	
Channel catfish	0.30 (0.12)	0.31 (0.13)	0.25 (0.16)
Northern pike	0.14 (0.10)	0.15 (0.11)	0.13 (0.13)
White bass	1.56 (0.75)	1.70 (0.83)	0.25 (0.25)
Yellow bass	0.78 (0.41)	0.86 (0.46)	
Rock bass	0.23	0.11	1.40

Table 6.3 continued.

Common name	ALL	Backwater, contiguous, shoreline	Impounded, shoreline
	(0.08)	(0.07)	(0.54)
Pumpkinseed	7.95	6.88	18.06
	(2.51)	(2.66)	(7.69)
Warmouth	0.05	0.05	
	(0.05)	(0.05)	
Orangespotted sunfish	0.01		0.13
	(0.01)		(0.13)
Bluegill	14.04	15.04	4.58
	(3.14)	(3.48)	(1.82)
Largemouth bass	1.79	1.63	3.32
	(0.48)	(0.49)	(1.77)
White crappie	1.50	1.63	0.26
	(0.72)	(0.79)	(0.26)
Black crappie	2.72	2.88	1.28
	(1.21)	(1.34)	(0.69)
Yellow perch	3.68	4.03	0.38
	(2.11)	(2.33)	(0.27)
Sauger	0.04	0.05	
	(0.04)	(0.05)	
Walleye	0.09	0.10	
	(0.07)	(0.07)	
Freshwater drum	6.34	6.49	4.89
	(2.48)	(2.74)	(2.90)